COUNTY OF LOS ANGELES – DEPARTMENT OF MENTAL HEALTH PROGRAM SUPPORT BUREAU – MHSA IMPLEMENTATION AND OUTCOMES DIVISION





WELLNESS • RECOVERY • RESILIENCE

Prevention & Early Intervention: Crisis Oriented Recovery Services (CORS)

Countywide Aggregate Practice Outcomes Dashboard Report Outcome Data Submission through February 22, 2016

Participating Legal Entities Include:

3 3		
ALCOTT CENTER FOR MHS	KEDREN COMMUNITY MENTAL HEALTH CTR	AUGUSTUS F HAWKINS FAMILY
ALMA FAMILY SERVICES	LOS ANGELES CHILD GUIDANCE	COASTAL API FAMILY MHC
AMANECER COMMUNITY COUNSELING SVC	MARYVALE	COMPTON FAMILY MHS
CALIF INSTITUTE HEALTH SOCIAL SVC	PERSONAL INVOLVEMENT CENTER INC	DMH AT HARBOR-UCLA MEDICAL
CENTER FOR INTEGRATED FAMILY HEALTH	SAINT JOHNS HEALTH CENTER	DOWNTOWN MENTAL HEALTH
CHILD AND FAMILY CENTER	SAN FERNANDO VALLEY COMMUNITY MHC	HOLLYWOOD MENTAL HEALTH
CHILD AND FAMILY GUIDANCE CENTER	SHIELDS FOR FAMILIES	LONG BEACH MHS ADULT CLIN
D VEAL FAMILY AND YOUTH SVC	SPECIAL SERVICE FOR GROUPS	NORTHEAST MENTAL HEALTH
DIDI HIRSCH PSYCHIATRIC SERVICE	ST JOSEPH CENTER	RIO HONDO COMMUNITY MHC
DIGNITY HEALTH DBA CAL HOSP MED CTR	STAR VIEW BEHAVIORAL HEALTH INC	ROYBAL FAMILY MHS
DREW CHILD DEVELOPMENT CORPORATION	THE VILLAGE FAMILY SERVICES	SAN ANTONIO MENTAL HEALTH
EL CENTRO DE AMISTAD INC	VIP COMMUNITY MENTAL HEALTH CTR INC	SAN PEDRO MENTAL HEALTH
ENKI HEALTH AND RESEARCH SYSTEMS I	L.A. COUNTY DMH	SOUTH BAY MHS
FOOTHILL FAMILY SERVICE	ANTELOPE VALLEY MHC	WEST CENTRAL FAMILY MHS
HERITAGE CLINIC	AMERICAN INDIAN COUNSELIN	
HILLVIEW MENTAL HEALTH CENTER INC	ARCADIA MHS	

Agencies submitting outcomes that are not approved to provide CORS by PEI Administration:

<u> </u>	<u> </u>
JEWISH FAMILY SERVICE OF LOSANGELES	L.A. COUNTY DMH
PENNY LANE CENTERS	ROYBAL SCHOOL BASED PROGRAM
TESSIE CLEVELAND COMM SRVC CORP	

Table 1. COF	Table 1. CORS Status Since Inception to February 22, 2016								
# of Clients Claimed to Practice	# of Clients Entered into PEI OMA	# of Tx Cycles in PEI OMA	Clients with Multiple Tx Cycles	Clients Completing Tx	Clients Dropping- Out of Tx	Clients Still -In Tx			
9195	30.77%	2930	3.25%	39.22%	26.86%	33.92%			
n=	2829	n=	92	1149	787	994			

Note 1: Clients Claimed was based on CORS being selected as the EBP in a PEI Plan and having ≥ 2 core services claimed to the practice starting July 1, 2011.

Note 2: Number of clients Completing Tx or Dropping-Out of Tx was determined by whether the EBP was said to be completed (e.g. answered "yes" or "no") in the PEI OMA.

Table 2. C	Table 2. Client Demographics - Clients Who Entered CORS											
	Age		Gender				Ethnicity	,		Prim	ary Langu	age
Total Number of Clients	Average	Female	Male	Unknown	African-American	Asian / Pacific Islander	Caucasian	Hispanic / Latino	Other	English	Spanish	Other
2829	19	52.03%	47.90%	0.07%	31.07%	1.84%	5.80%	56.24%	5.05%	70.34%	27.39%	2.26%
	n=	1472	1355	2	879	52	164	1591	143	1990	775	64

Note 1: Age is calculated at the date of the first EBP.

Note 2: Percentages may not total 100 due to missing data and/or rounding errors.

Table 3. Top 5 Most Frequently Reported DSM-IV Primary Axis Diagnosis - Clients Who Entered
CORS

Total Treatment Cycles	Depressive Disorder NOS	Mood Disorder NOS	Attention- Deficit/Hyperactivity Disorder, Combined Type or Hyperactive Impulse Type	Disruptive Behavior Disorder NOS	Anxiety Disorder NOS	Other
2930	14.74%	9.62%	6.45%	5.53%	5.26%	54.06%
n=	432	282	189	162	154	1584

NOTE: The above table reflects diagnoses entered PEI OMA from July 1, 2011 through October 27, 2015.

*See breakout of "Other" on the supplemental information page.

Table 4. Program Process Data - Clients Who Entered CORS							
Outcome Measures Administered	Pre-Test with Scores	Post-test with Scores	Clients Who Completed both a Pre and Post Measure with Scores				
Youth Outcome Questionnaire - 2.01 (Parent)	84.23%	63.95%	31.18%				
n=	1645	690	609				
Ackn=	1953	1079	1953				
Youth Outcome Questionnaire – Self Report – 2.0 (YOQ- SR)	82.16%	64.60%	27.52%				
n=	806	312	270				
Ackn=	981	483	981				
Outcome Questionnaire - 45.2	82.63%	51.02%	19.46%				
n=	709	200	167				
Ackn=	858	392	858				

Note 1: Number of acknowledged measures (Ackn=) is determined by the number of required measures that receive a score or an unable to collect reason code.

Note 2: The % indicated for Pre-test with scores, Post-test with scores, and both a Pre- and Post-test with scores is calculated by dividing the (n=#) by the number acknowledged (Ackn=#) in the PEI OMA system for each measure. The number acknowledged (Ackn=#) for those with Pre and Post scores is an estimate based on the greatest number of matches that could be expected given the number of Pre scores acknowledged.

Table 5a	able 5a. Top Reasons Given for "Unable to Collect"							
Questionnaire - 2.01 arent)	Total Pre 308	Parent/care provider unavailable	Administration date exceeds acceptable range	Outcome measure unavailable	Invalid outcome measure	Parent/care provider refused	Other Reasons	
tionr (percent	36.36%	25.65%	11.04%	10.39%	6.17%	10.39%	
e Questi (Parent)	n	112	79	34	32	19	32	
Outcome (P	Total Post 389	Parent/care provider unavailable	Administration date exceeds acceptable range	Premature termination	Outcome measure unavailable	Parent/care provider refused	Other Reasons	
Youth	percent	37.02%	17.22%	16.20%	10.80%	6.17%	12.60%	
	n	144	67	63	42	24	49	

Table 5b	Table 5b. Top Reasons Given for "Unable to Collect"								
Questionnaire 2.0 (YOQ-SR)	Total Pre 175	Administration date exceeds acceptable range	Outcome measure unavailable	Client unavailable	Invalid outcome measure	Client refused	Other Reasons		
lesti (YO	percent	24.00%	19.43%	18.29%	12.00%	10.29%	16.00%		
	n	42	34	32	21	18	28		
Youth Outcome Self Report -	Total Post 171	Client unavailable	Premature termination	Administration date exceeds acceptable range	Lost contact with client	Outcome measure unavailable	Other Reasons		
	percent	22.81%	20.47%	18.71%	13.45%	8.77%	15.79%		
	n	39	35	32	23	15	27		

Table 50	Table 5c. Top Reasons Given for "Unable to Collect"								
- 45.2	Total Pre 149	Outcome measure unavailable	Clinician not trained in outcome measure	Client refused	Administration date exceeds acceptable range	Client unavailable	Other Reasons		
naire	percent	18.12%	17.45%	15.44%	15.44%	14.09%	19.46%		
tion	n	27	26	23	23	21	29		
Outcome Questionnaire	Total Post 192	Client unavailable	Lost contact with client	Administration date exceeds acceptable range	Premature termination	Client refused	Other Reasons		
	percent	30.21%	21.35%	9.38%	9.38%	8.85%	20.83%		
	n	58	41	18	18	17	40		

Т	Table 6. Service Delivery Data – Clients Who Completed CORS							
Т	Total reatment Cycles	Average Length of Treatment in Weeks	-	ge of nt Weeks	Average Number of Sessions	Range of	f Sessions	
	1149	10	Min Max		8	Min	Max	
		10	0	69	J	1	105	

Note: Completed CORS is defined as having a 'yes' for completion indicated in the PEI OMA.

Table 7. Outcome Data* – Clients who Completed CORS								
		Percent Improvement	Percent of Clients Showing Reliable Change* from Pre- CORS to Post-CORS					
		from Pre to Post	Positive Change	No change	Negative Change			
Youth Outcome		27.400/	44.500/	45 650/	0.050/			
Questionnaire (YOQ) - 2.01	TOTAL	27.49%	44.50%	45.65%	9.85%			
(Parent)		(n=609)	271	278	60			
Youth Outcome								
Questionnaire	TOTAL	28.68%	40.74%	50.74%	8.52%			
- Self Report - 2.0 (YOQ-SR)		(n=270)	110	137	23			
Outcome								
Questionnaire	TOTAL	20.98%	38.32%	53.89%	7.78%			
- 45.2		(n=167)	64	90	13			

[‡]Please see Appendix A. for a description of the

CORS outcome measures and the outcome indicators (percent improvement in average scores; and, percent of clients showing reliable change).

Note 2: Possible YOQ-Parent Total Scores can range from -16 to 240, with a clinical cutpoint of 46


Note 3: Possible YOQ-SR Total Scores can range from -16 to 240, with a clinical cutpoint of 47.


Note 4: Possible OQ-45.2 Total Score can range from 0 to 180, with a clinical cutpoint of 64.


Note 5: Aggregate outcome data based on fewer than 20 clients are not reported.


Note 6: Positive Change indicates that the scores decreased from the pre to the post measures.























Appendix

Youth Outcomes Questionnaires (YOQ (Parent) and YOQ-SR)

The Youth Outcome Questionnaire is a 64-item parent-report that assesses global distress in a child's/adolescent's life from 4-17 years of age. The YOQ-SR is the Self-report version of the YOQ and is completed by the child/adolescent him or herself. Scores on both measures can range from -16 to 240. Scores of 46 or higher are most similar to a clinical population on the YOQ. A score of 47 is most similar to that of a clinical population on the YOQ-SR.

Outcomes Questionnaires (OQ)

The Outcome Questionnaire is a 45-item self-report that assesses global distress in a client's life from ages 19 and older. Total Scores on this measure can range from 0 to 180, with scores of 64 or higher indicating clinical significance.

Reliable Change Index

When comparing Pre and Post scores, it is very helpful to know whether the change reported represents the real effects of the treatment or errors in the system of measurement. The Reliability of Change Index (RCI) is a statistical way of helping to insure that the change recorded between pre and post assessments exceeds that which would be expected on the basis of measurement error alone. The RCI has been calculated using the Jacobson and Truax (1991) method and indicates when change exceeds that which would be expected on the basis of error at the p<.05 probability level. For a more in-depth discussion of Reliability of Change see Jacobson, N. S., & Truax. P. (1991). Clinical Significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology*, 59, 12-19. Also see Wise, E. A. (2004). Methods for analyzing psychotherapy outcomes: A review of clinical significance, reliable change, and recommendations for future directions. *Journal of Personality Assessment*, 82(1), 50-59.

The number and percent of clients experiencing positive change, no change and negative change are recorded in table 6. Healthful change in each of the measures cited here means that scores have <u>decreased</u> in value from pre to post test administrations (i.e. recorded a negative change on the RCI). To help avoid confusion, healthful reliable change is presented as positive while unhealthful reliable change is presented as negative change.

Supplemental Information for Table 3 "Other" Diagnosis Category

Supplemental information for Table 6 Other Biagnosis		
DSM-IV Primary Axis Diagnosis - Clients Who Entered CORS	n	Percent
Adjustment Disorder With Anxiety and Depressed Mood	140	6.29%
Post-Traumatic Stress Disorder	128	5.75%
Adjustment Disorder With Depressed Mood	125	5.61%
Adjustment Disorder Unspecified	92	4.13%
Psychotic Disorder NOS	91	4.09%
Oppositional Defiant Disorder	83	3.73%
Major Depressive Disorder, Recurrent, Moderate	64	2.87%
Adjustment Disorder W/Mixed Disturbance Emotion and Conduct	61	2.74%
Adjustment Disorder With Anxiety	53	2.38%
Major Depressive Disorder, Rec., Severe W/O Psychotic Features	53	2.38%
Major Depressive Disorder, Rec., Severe With Psychotic Features	53	2.38%
Dysthymic Disorder	49	2.20%
Generalized Anxiety Disorder	48	2.16%
Attention-Deficit/Hyperactivity Disorder NOS	47	2.11%
Major Depressive Disorder, Single Episode, Unspecified	44	1.98%
Disorder of Infancy, Childhood, or Adolescence NOS	39	1.75%
Attention-Deficit/Hyperactivity Disorder, Predominantly Inattentive Type	38	1.71%
Adjustment Disorder With Disturbance of Conduct	35	1.57%
Major Depressive Disorder, Single Episode, Moderate	28	1.26%
Major Depressive Disorder, Single Episode, Mild	26	1.17%
Major Depressive Disorder, Recurrent, Unspecified	19	0.85%
Bipolar Disorder NOS	19	0.85%
Acute Stress Disorder	19	0.85%
Major Depressive Disorder, Recurrent, Mild	16	0.72%
Schizophrenia, Paranoid Type	15	0.67%
Schizoaffective Disorder	14	0.63%
Major Depressive Disorder, SE, Severe W/O Psychotic Features	14	0.63%
Separation Anxiety Disorder	14	0.63%
Panic Disorder Without Agoraphobia	13	0.58%
Pervasive Developmental Disorder NOS/Asperger's Disorder/Rett's Disorder	12	0.54%
Diagnosis Deferred (code invalid as Secondary/Axis II Diagnosis)	12	0.54%
Conduct Disorder, Adolescent-Onset Type	11	0.49%
Obsessive-Compulsive Disorder	10	0.45%
Major Depressive Disorder, SE, Severe With Psychotic Features	10	0.45%
Impulse-Control Disorder NOS	9	0.40%
Panic Disorder With Agoraphobia	8	0.36%
Mood Disorder Due to(Indicate Medical Condition)	7	0.31%
Conduct Disorder, Childhood-Onset Type	6	0.27%
• • • • • • • • • • • • • • • • • • • •		

TOTAL	1585	71.17%
Bipolar I Disorder, Single Manic Episode, Moderate	1	0.04%
Bereavement	1	0.04%
Occupational Problem	1	0.04%
Psychotic Disorder Due to(Indicate Medical Condition) with Delusions	1	0.04%
Drug Related Disorder NOS	1	0.04%
Drug Induced Mood Disorder	1	0.04%
Drug Induced Psychotic Disorder, With Hallucinations	1	0.04%
Drug Induced Psychotic Disorder, With Delusions	1	0.04%
Bipolar I, SME, Severe W/O Psychotic Features	1	0.04%
Bipolar I Disorder, Single Manic Episode, Moderate	1	0.04%
Bipolar I Disorder, Single Manic Episode, Mild	1	0.04%
Delusional Disorder	1	0.04%
Autistic Disorder	1	0.04%
Bipolar I Disorder, MRE, Manic, Severe With Psychotic Features	1	0.04%
Major Depressive Disorder, Recurrent, In Full Remission	1	0.04%
Major Depressive Disorder, Recurrent, In Partial Remission	1	0.04%
Bipolar I Disorder, MRE, Mixed, Severe W/O Psychotic Features	1	0.04%
Bipolar I Disorder, MRE, Unspecified	1	0.04%
Dissociative Amnesia	1	0.04%
Conversion Disorder	1	0.04%
Specific Phobia	1	0.04%
Polysubstance Dependence	1	0.04%
Unspecified Mental Disorder (Nonpsychotic)	1	0.04%
Bulemia Nervosa	1	0.04%
Intermittent Explosive Disorder	1	0.04%
Selective Mutism	1	0.04%
Identity Problem	1	0.04%
Encopresis W/Constipation and Overflow Incontinence	1	0.04%
Learning Disorder NOS	2	0.09%
Cannabis Abuse	2	0.09%
Bipolar I Disorder, MRE, Mixed, Severe With Psychotic Features	2	0.09%
Bipolar I Disorder, MRE, Depressed, Severe With Psychotic Features	2	0.09%
Schizophrenia, Undifferentiated Type	2	0.09%
Bipolar I Disorder, Single Manic Episode, Unspecified	3	0.13%
Social Phobia	4	0.18%
Brief Psychotic Disorder	4	0.18%
Bipolar II Disorder	5	0.22%
No Diagnosis (code invalid as Secondary/Axis III Diagnosis)	6	0.27%

^{*}Percentage is not exact due to rounding errors.